


Prenatal Development and the Newborn

45-2 What is the course of prenatal development, and how do teratogens affect that development?


Conception

Nothing is more natural than a species reproducing itself. And nothing is more wondrous. With humans, the process starts when a woman's ovary releases a mature egg—a cell roughly the size of the period at the end of this sentence. The woman was born with all the immature eggs she would ever have, although only 1 in 5000 will ever mature and be released. A man, in contrast, begins producing sperm cells at puberty. For the rest of his life, 24 hours a day, he will be a nonstop sperm factory, with the rate of production—in the beginning more than 1000 sperm during the second it takes to read this phrase—slowing with age.

Like space voyagers approaching a huge planet, the 200 million or more deposited sperm begin their race upstream, approaching a cell 85,000 times their own size. The relatively few reaching the egg release digestive enzymes that eat away its protective coating (**FIGURE 45.2a**). As soon as one sperm penetrates that coating and is welcomed in (Figure 45.2b), the egg's surface blocks out the others. Before half a day elapses, the egg nucleus and the sperm nucleus fuse. The two have become one. Consider it your



(a)



(b)

Figure 45.2

Life is sexually transmitted (a) Sperm cells surround an egg. (b) As one sperm penetrates the egg's jellylike outer coating, a series of chemical events begins that will cause sperm and egg to fuse into a single cell. If all goes well, that cell will subdivide again and again to emerge 9 months later as a 100-trillion-cell human being.

most fortunate of moments. Among 200 million sperm, the one needed to make you, in combination with that one particular egg, won the race. And so it was for innumerable generations before us. If any one of our ancestors had been conceived with a different sperm or egg, or died before conceiving, or not chanced to meet the partner or . . . the mind boggles at the improbable, unbroken chain of events that produced you and me.

Prenatal Development

Fewer than half of all fertilized eggs, called **zygotes**, survive beyond the first 2 weeks (Grobstein, 1979; Hall, 2004). But for you and me, good fortune prevailed. One cell became 2, then 4—each just like the first—until this cell division had produced some 100 identical cells within the first week. Then the cells began to differentiate—to specialize in structure and function. How identical cells do this—as if one decides “I’ll become a brain, you become intestines!”—is a puzzle that scientists are just beginning to solve.

About 10 days after conception, the zygote attaches to the mother’s uterine wall, beginning approximately 37 weeks of the closest human relationship. The zygote’s inner cells become the **embryo** (FIGURE 45.3a). The outer cells become the *placenta*, the life-link that transfers nutrients and oxygen from mother to embryo. A healthy and well-nourished mother helps form a healthy baby-to-be. Over the next 6 weeks, the embryo’s organs begin to form and function. The heart begins to beat.

For 1 in 270 sets of parents, though, there is a bonus. Two heartbeats will reveal that the zygote, during its early days of development, has split into two. If all goes well, two genetically identical babies will start life together some 8 months later (Module 14).

By 9 weeks after conception, an embryo looks unmistakably human (Figure 45.3b). It is now a **fetus** (Latin for “offspring” or “young one”). During the sixth month, organs such as the stomach have developed enough to give the fetus a good chance of survival if born prematurely.

At each prenatal stage, genetic and environmental factors affect our development. By the sixth month, microphone readings taken inside the uterus reveal that the fetus is responsive to sound and is exposed to the sound of its mother’s muffled voice (Ecklund-Flores, 1992; Hepper, 2005). Immediately after birth, emerging from living 38 or so weeks underwater, newborns prefer her voice to another woman’s or to their father’s (Busnel et al., 1992; DeCasper et al., 1984, 1986, 1994). They also prefer hearing their mother’s language. If she spoke two languages during pregnancy, they display interest in both (Byers-Heinlein et al., 2010). And just after birth, the melodic ups and downs of newborns’ cries bear the tuneful signature of their mother’s native tongue (Mampe et al., 2009). Babies born

zygote the fertilized egg; it enters a 2-week period of rapid cell division and develops into an embryo.

embryo the developing human organism from about 2 weeks after fertilization through the second month.


fetus the developing human organism from 9 weeks after conception to birth.

Figure 45.3


Prenatal development (a) The embryo grows and develops rapidly. At 40 days, the spine is visible and the arms and legs are beginning to grow. (b) By the end of the second month, when the fetal period begins, facial features, hands, and feet have formed. (c) As the fetus enters the fourth month, its 3 ounces could fit in the palm of your hand.



(a)



(b)



(c)

Anatomical Travelogue/Science Source

Gary Retherford/Science Source

Petit Format/Science Source

to French-speaking mothers tend to cry with the rising intonation of French; babies born to German-speaking mothers cry with the falling tones of German. Would you have guessed? The learning of language begins in the womb.

In the 2 months before birth, fetuses demonstrate learning in other ways, as when they adapt to a vibrating, honking device placed on their mother’s abdomen (Dirix et al., 2009). Like people who adapt to the sound of trains in their neighborhood, fetuses get used to the honking. Moreover, 4 weeks later, they recall the sound (as evidenced by their biased response, compared with reactions of those not previously exposed).

Sounds are not the only stimuli fetuses are exposed to in the womb. In addition to transferring nutrients and oxygen from mother to fetus, the placenta screens out many harmful substances, but some slip by. **Teratogens**, agents such as viruses and drugs, can damage an embryo or fetus. This is one reason pregnant women are advised not to drink alcoholic beverages. A pregnant woman never drinks alone. As alcohol enters her bloodstream, and her fetus’, it depresses activity in both their central nervous systems. Alcohol use during pregnancy may prime the woman’s offspring to like alcohol and may put them at risk for heavy drinking and alcohol use disorder during their teens. In experiments, when pregnant rats drank alcohol, their young offspring later displayed a liking for alcohol’s taste and odor (Youngentob et al., 2007, 2009).

Even light drinking or occasional binge drinking can affect the fetal brain (Braun, 1996; Ikonomidou et al., 2000; Sayal et al., 2009). Persistent heavy drinking puts the fetus at risk for birth defects and for future behavior problems, hyperactivity, and lower intelligence. For 1 in about 800 infants, the effects are visible as **fetal alcohol syndrome (FAS)**, marked by lifelong physical and mental brain abnormalities (May & Gossage, 2001). The fetal damage may occur because alcohol has an *epigenetic effect*: It leaves chemical marks on DNA that switch genes abnormally on or off (Liu et al., 2009).

The Competent Newborn

45.3 What are some newborn abilities, and how do researchers explore infants’ mental abilities?

Babies come with software preloaded on their neural hard drives. Having survived prenatal hazards, we as newborns came equipped with automatic reflex responses ideally suited for our survival. We withdrew our limbs to escape pain. If a cloth over our face interfered with our breathing, we turned our head from side to side and swiped at it.

New parents are often in awe of the coordinated sequence of reflexes by which their baby gets food. Thanks to the *rooting reflex*, when something touches their cheek, babies turn toward that touch, open their mouth, and vigorously root for a nipple. Finding one, they automatically close on it and begin *sucking*—which itself requires a coordinated sequence of reflexive *tonguing*, *swallowing*, and *breathing*. Failing to find satisfaction, the hungry baby may cry—a behavior parents find highly unpleasant and very rewarding to relieve.



Lofzapsas Photography, Inc./Corbis



Aasa Images/Getty Images

Prepared to feed and eat Animals are predisposed to respond to their offspring’s cries for nourishment.

FYI

Prenatal development

zygote: conception to 2 weeks
embryo: 2 to 9 weeks
fetus: 9 weeks to birth

“You shall conceive and bear a son. So then drink no wine or strong drink.” —JUDGES 13:7

teratogens (literally, “monster maker”) agents, such as chemicals and viruses, that can reach the embryo or fetus during prenatal development and cause harm.

fetal alcohol syndrome (FAS) physical and cognitive abnormalities in children caused by a pregnant woman’s heavy drinking. In severe cases, signs include a small, out-of-proportion head and abnormal facial features.

“I felt like a man trapped in a woman’s body. Then I was born.” —COMEDIAN CHRIS BLISS

habituation decreasing responsiveness with repeated stimulation. As infants gain familiarity with repeated exposure to a visual stimulus, their interest wanes and they look away sooner.


The pioneering American psychologist William James presumed that the newborn experiences a “blooming, buzzing confusion,” an assumption few people challenged until the 1960s. But then scientists discovered that babies can tell you a lot—if you know how to ask. To ask, you must capitalize on what babies can do—gaze, suck, turn their heads. So, equipped with eye-tracking machines and pacifiers wired to electronic gear, researchers set out to answer parents’ age-old questions: What can my baby see, hear, smell, and think?

Consider how researchers exploit **habituation**—a decrease in responding with repeated stimulation. We saw this earlier when fetuses adapted to a vibrating, honking device placed on their mother’s abdomen. The novel stimulus gets attention when first presented. With repetition, the response weakens. This seeming boredom with familiar stimuli gives us a way to ask infants what they see and remember.

An example: Researchers have used *visual preference* to “ask” 4-month-olds how they recognize cats and dogs (Quinn, 2002; Spencer et al., 1997). First, they showed the infants a series of images of either cats or dogs. Then they showed them hybrid cat-dog images (**FIGURE 45.4**). Which of those two animals do you think the infants would find more novel (measured in looking time) after seeing a series of cats? It was the hybrid animal with the dog’s head (and vice versa if they previously viewed dogs). This suggests that infants, like adults, focus first on the face, not the body.

Figure 45.4

Quick—which is the cat? Researchers used cat-dog hybrid images such as these to test how infants categorize animals.




Indeed, even as newborns, we prefer sights and sounds that facilitate social responsiveness. We turn our heads in the direction of human voices. We gaze longer at a drawing of a face-like image (**FIGURE 45.5**). We prefer to look at objects 8 to 12 inches away. Wonder of wonders, that just happens to be the approximate distance between a nursing infant’s eyes and its mother’s (Maurer & Maurer, 1988).

Within days after birth, our brain’s neural networks were stamped with the smell of our mother’s body. Week-old nursing babies, placed between a gauze pad from their mother’s bra and one from another nursing mother, have usually turned toward the smell of their own mother’s pad (MacFarlane, 1978). What’s more, that smell preference lasts. One experiment capitalized on the fact that some nursing mothers in a French maternity ward applied a

Figure 45.5

Newborns’ preference for faces When shown these two stimuli with the same elements, Italian newborns spent nearly twice as many seconds looking at the face-like image (Johnson & Morton, 1991). Canadian newborns display the same apparently inborn preference to look toward faces (Mondloch et al., 1999).



balm with a chamomile scent to prevent nipple soreness (Delaunay-El Allam, et al., 2010). Twenty-one months later, their toddlers preferred playing with chamomile-scented toys! Their peers who had not sniffed the scent while breast feeding showed no such preference. (This makes one wonder: Will adults who as babies associated chamomile scent with their mother’s breast become devoted chamomile tea drinkers?)

Before You Move On

▶ ASK YOURSELF

Are you surprised by the news of infants’ competencies? Remember hindsight bias from Module 4? Is this one of those cases where it feels like you “knew it all along”?

▶ TEST YOURSELF

Your friend’s older sister—a regular drinker—hopes to become pregnant soon and has stopped drinking. Why is this a good idea? What negative effects might alcohol consumed during pregnancy have on a developing fetus?

Answers to the Test Yourself questions can be found in Appendix E at the end of the book.

Module 45 Review

45-1

What three issues have engaged developmental psychologists?

- *Developmental psychologists* study physical, mental, and social changes throughout the life span.
- They focus on three issues: nature and nurture (the interaction between our genetic inheritance and our experiences); continuity and stages (whether development is gradual and continuous or a series of relatively abrupt changes); and stability and change (whether our traits endure or change as we age).

45-3

What are some newborn abilities, and how do researchers explore infants’ mental abilities?

- Babies are born with sensory equipment and reflexes that facilitate their survival and their social interactions with adults. For example, they quickly learn to discriminate their mother’s smell and sound.
- Researchers use techniques that test *habituation*, such as the visual-preference procedure, to explore infants’ abilities.

45-8

What is the course of prenatal development, and how do teratogens affect that development?

- The life cycle begins at conception, when one sperm cell unites with an egg to form a *zygote*.
- The *zygote*’s inner cells become the *embryo*, and in the next 6 weeks, body organs begin to form and function.
- By 9 weeks, the *fetus* is recognizably human.
- *Teratogens* are potentially harmful agents that can pass through the placental screen and harm the developing embryo or fetus, as happens with *fetal alcohol syndrome*.